

## 20. Literaturverzeichnis

- Aaronson NK, Ahmedzai S, Bergman B, et al. The European Organization for Research and Treatment of Cancer QLQ-C30: a quality-of-life instrument for use in international clinical trials in oncology. *J Natl Cancer Inst* 1993; 85(5): 365-376
- Abrey LE, Rosenblum MK, Papadopoulos E, et al. High dose chemotherapy with autologous stem cell rescue in adults with malignant primary brain tumors. *J Neurooncol* 1999; 44 (2): 147-153
- Adamson PC, Balis FM, Arndt CA, et al. Intrathecal 6-mercaptopurine: preclinical pharmacology, phase I/II trial, and pharmacokinetic study. *Cancer Res* 1991; 51 (22): 6079-6083
- Aggarwal R, Yeung D, Kumar P, et al. Efficacy and feasibility of stereotactic radio-surgery in the primary management of unfavorable pediatric ependymoma. *Radiother Oncol* 1997; 43: 269-273
- Akay KM, Izci Y, Baysefer A, et al. Surgical outcomes of cerebellar tumors in children. *Pediatr Neurosurg* 2004; 40(5): 220-225
- Allen JC, Helson L. High-dose cyclophosphamide chemotherapy for recurrent CNS tumors in children. *J Neurosurg* 1981; 55: 749-756
- Allen JC, Epstein F. Medulloblastoma and other primary malignant tumors of the CNS. *J Neurosurg* 1982; 57: 446-452
- Allen JC, Walker R, Luks E, et al. Carboplatin and recurrent childhood brain tumors. *J Clin Oncol* 1987; 5: 459-463
- Balon HR, Goldsmith SJ, Siegel BA, et al. Society of Nuclear Medicine Procedure Guideline for Somatostatin Receptor Scintigraphy with In-111 Pentetetrotide; version 1.0, approved February 21<sup>st</sup>, 2001.
- Balter-Seri J, Mor C, Shuper A, et al. Cure of recurrent medulloblastoma: the contribution of surgical resection at relapse. *Cancer* 1997; 79 (6): 1241-1247
- Becker LE, Hinton D. Primitive neuroectodermal tumors of the central nervous system. *Human Pathol* 1983; 14: 538-550
- Belza MG, Donaldson SS, Steinberg GK, et al. Medulloblastoma: freedom from relapse longer than 8 years - a therapeutic cure? *J Neurosurg* 1991; 75 (4): 575-582
- Berg SL, Balis FM, Zimm S, et al. Phase I/II trial and pharmacokinetics of intrathecal diaziquone in refractory meningeal malignancies. *J Clin Oncol* 1992; 10 (1): 143-148
- Berry MP, Jenkin RD, Colin MB, et al. Radiation treatment for medulloblastoma. A 21-year review. *J Neurosurg* 1981; 55: 43-51
- Blaney SM, Poplack DG. New cytotoxic drugs for intrathecal administration. *J Neurooncol* 1998; 38: 219-223
- Bleyer WA, Poplack DG, Simon RM. "Concentration x time" methotrexate via a subcutaneous reservoir: a less toxic regimen for intraventricular chemotherapy of central nervous system neoplasms. *Blood* 1978; 51: 835-842
- Bloom HJG. The role of adjuvant chemotherapy in the primary treatment of medulloblastoma and ependymoma. In: Pochedly C (ed.). *Controversies in pediatric and adolescent hematology and oncology*. CV Mosby, St. Louis, 1989
- B. Borgelt et al. *Int J Radiat Oncol Biol Phys* 1980; 6: 1-9 (6)). The palliation of brain metastases: final results of the first two studies by the Radiation Therapy Oncology Group.
- Broniscer A, Nicolaides TP, Dunkel IJ, et al. High-dose chemotherapy with autologous stem-cell rescue in the treatment of patients with recurrent non-cerebellar primitive neuroectodermal tumors. *Pediatr Blood Cancer* 2004; 42 (3): 261-267
- Busca A, Miniero R, Besenzi L, et al. Etoposide-containing regimens with autologous bone marrow transplantation in children with malignant brain tumors. *Childs Nerv Syst* 1997; 13 (11-12): 572-577
- Calaminus G, Ittner E, Neubauer M, et al. Prospective evaluation of late effects and quality of life (QOL) in patients with ALL and brain tumors in Germany. *Med Pediatr Oncol* 2002; 39: 345
- Calaminus G, Weinsbach S, Teske C, Göbel U. Quality of life in children and adolescents with cancer. First results of an evaluation of 49 patients with the PEDQOL questionnaire. *Klin Pädiatr* 2000; 212(4): 211-215
- Castello MA, Clerico A, Deb G, et al. High-dose carboplatin in combination with etoposide (JET regimen) for childhood brain tumors. *Am J Pediatr Hematol Oncol* 1990; 12 (3): 297-300
- Cereda C, Calaminus G, Varotto S, et al. Searching for a tool to assess quality of life (QOL) in childhood cancer survivors – results of the PEDQOL in a cohort of Italian children and adolescents. *Med Pediatr Oncol* 2003; 41: 271

- Cinatl J, Kotchetkov R, Blaheta R, et al., Induction of differentiation and suppression of malignant phenotype of human neuroblastoma BE(2)-C cells by valproic acid: enhancement by combination with interferon-alpha. *Int J Oncol* 2002; 20 (1): 97-106
- Chamberlain MC, Prados MD, Silver P, et al. A phase II trial of oral melphalan in recurrent primary brain tumors. *Am J Clin Oncol* 1988; 11 (1): 52-54
- Chamberlain MC, Kormanik PA. Chronic oral VP-16 for recurrent medulloblastoma. *Pediatr Neurol* 1997a; 17 (3): 230-234
- Chamberlain MC. Pediatric leptomeningeal metastases: outcome following combined therapy. *J Child Neurol* 1997b; 12: 53-59
- Chamberlain MC. Radioisotope CSF flow studies in leptomeningeal metastases. *J Neurooncology* 1998; 38: 135-140
- Chamberlain MC. Recurrent intracranial ependymoma in children: salvage therapy with oral etoposide. *Pediatr Neurol* 2001; 24 (2): 117-121
- Chang CH, Housepian EM, Herbert C Jr. An operative staging system and a megavoltage radiotherapeutic technic for cerebellar medulloblastomas. *Radiology* 1969; 93: 1351-359
- Cohen, BH, Packer RJ. Chemotherapy for medulloblastomas and primitive neuroectodermal tumors. *J Neurooncol* 1996; 29: 55-68
- Dallorso S, Manzitti C, Morreale G, et al. High dose therapy and autologous hematopoietic stem cell transplantation in poor risk solid tumors of childhood. *Haematologica* 2000; 85 (11 Suppl): 66-70
- Das A, Banik NL, Patel SJ, Ray SK. Dexamethasone protected human glioblastoma U87MG cells from temozolomide induced apoptosis by maintaining Bax:Bcl-2 ratio and preventing proteolytic activities. *Mol Cancer* 2004; 3(1): 36
- Däumling E, Dierksen A, Wolff J, Jürgens H: FMH, a new instrument to measure quality of life. *Klin Pädiatr* 1994; 206: 351
- Davidson A, Gowing R, Lowis S, et al. Phase II study of 21 day schedule oral etoposide in children. New Agents Group of the United Kingdom Children's Cancer Study Group (UKCCSG). *Eur J Cancer* 1997; 33 (11): 1816-1822
- DeAngelis LM. Current diagnosis and treatment of leptomeningeal metastasis. *J Neurooncol* 1998; 38: 245-252
- Dunbar SF, Tarbell NJ, Kooy HM, Alexander E 3rd, Black PM, Barnes PD, Goumnerova L, Scott RM, Pomeroy SL, LaVally B. Stereotactic radiotherapy for pediatric and adult brain tumors: preliminary report. *Int J Radiat Oncol Biol Phys* 1994; 30 (3): 531-539
- Duffner PK, Cohen ME. Extraneural metastases in childhood brain tumors. *Ann Neurol* 1981; 10: 261-265
- Duffner PK, Horowitz ME, Krischer JP, et al. Postoperative chemotherapy and delayed radiation in children less than three years of age with malignant brain tumors. *New Engl J Med* 1993; 328: 1725-1731
- Dunkel IJ, Finlay JL. High dose chemotherapy with autologous stem cell rescue for patients with medulloblastoma. *J Neurooncol* 1996; 29: 69-74
- Dunkel IJ, Boyett JM, Yates A, et al. High-dose carboplatin, thiotepa, and etoposide with autologous stem-cell rescue for patients with recurrent medulloblastoma. Children's Cancer Group. *J Clin Oncol* 1998; 16: 222-228
- Dupuis-Girod S, Hartmann O, Benhamou E, et al. Will high dose chemotherapy followed by autologous bone marrow transplantation supplant crano-spinal irradiation in young children treated for medulloblastoma? *J Neurooncol* 1996; 27: 87-98
- Evans AE, Jenkin RDT, Sposto R, et al. The treatment of medulloblastoma: Results of a prospective randomized trial of radiation therapy with and without CCNU, vincristine, and prednisone. *J Neurosurg* 1990; 72: 572-582
- Farina P, Marzillo G, D'Incalci M. High-performance liquid chromatography determination of 4'-demethyl-epipodophyllotoxin-9-(4,6-O-ethylidene beta-D-glucopyranoside) (VP 16-213) in human plasma. *J Chromatogr* 1981; 222: 141-145
- Fazekas F. Magnetic resonance signal abnormalities in asymptomatic individuals: their incidence and functional correlates. *Eur Neurol* 1989; 29 (3):164-168
- Finlay JL. The role of high-dose chemotherapy and stem cell rescue in treatment of malignant brain tumors: a reappraisal. *Pediatr Transplant* 1999; 3 Suppl: 87-95
- Finlay, J. L., S. Goldman, M. C. Wong, et al. Pilot study of high-dose thiotepa and etoposide with autologous bone marrow rescue in children and young adults with recurrent CNS tumors. The Children's Cancer Group. *J Clin Oncol* 1996; 14: 2495-2503

- Fleischhack G, Reif S, Hasan C, et al. Feasibility of intraventricular administration of etoposide in patients with metastatic brain tumors. *Br J Cancer* 2001; 84: 1453-1459
- Fröhwald MC, Rickert CH, O'Dorisio MS, et al. Somatostatin receptor subtype 2 is expressed by supratentorial primitive neuroectodermal tumors of childhood and can be targeted for somatostatin receptor imaging. *Clin Cancer Res* 2004; 10 (9): 2997-3006
- Gajjar A, Kühl J, Epelman S, et al. Chemotherapy of medulloblastoma. *Childs Nerv Syst* 1999; 15 (10): 554-562
- Gardner S, Baker D, Belasco J, et al. Phase I dose escalation of temozolamide with thiotepa and carboplatin with autologous stem cell rescue (ASCR) in patients with recurrent/refractory central nervous system (CNS) tumors. 11<sup>th</sup> International Symposium on Pediatric Neuro-Oncology, Boston, 13<sup>th</sup>-16<sup>th</sup> June 2004 (abstr)
- Gaze MN, Smith DB, Rampling RP, et al. Combination chemotherapy for primitive neuroectodermal and other malignant brain tumours. *Clin Oncol (R Coll Radiol)* 1994; 6 (2): 110-115
- Gentet JC, Doz F, Bouffet E, et al. Carboplatin and VP-16 in medulloblastoma: A phase II study of the French Study of Pediatric Oncology (SFOP). *Med Ped Oncol* 1994; 23: 422-427
- Geyer JR, Zeltzer PM, Boyett JM, et al. Survival of infants with primitive neuroectodermal tumors or malignant ependymomas of the CNS treated with eight drugs in 1 day: A report from the Childrens Cancer Group. *J Clin Oncol* 1994; 12: 1607-1615
- Goldwein JR, Leahy JM, Packer RJ, et al. Intracranial ependymomas in children. *Int J Radiat Oncol Biol Phys* 1990; 19: 1497-1502
- Goodman R. The extended version of the Strengths and Difficulties Questionnaire as a guide to child psychiatric caseness and consequent burden. *J Child Psychol Psychiatry* 1999; 40(5): 791-800
- Grabb PA, Lunsford LD, Albright AL, et al. Stereotactic radiosurgery for glial neoplasms of childhood. *Neurosurgery* 1996; 38 (4): 696-701; discussion 701-702
- Graham ML, Herndon JE, Casey JR, et al. High-dose chemotherapy with autologous stem-cell rescue in patients with recurrent and high-risk pediatric brain tumors. *J Clin Oncol* 1997; 15: 1814-1823
- Grill J, Kalifa C, Doz F, et al. A high-dose busulfan-thiotepa combination followed by autologous bone marrow transplantation in childhood recurrent ependymoma. A phase-II study. *Pediatr Neurosurg* 1996; 25: 7-12
- Groll AH, Walsh TJ. Antifungal Chemotherapy: Recent advances and current perspectives. *Swiss Medical Weekly* 2002; 132: 303-311
- Gururangan S, McLaughlin C, Quinn J, et al. High-dose chemotherapy with autologous stem-cell rescue in children and adults with newly diagnosed pineoblastomas. *J Clin Oncol* 2003; 21 (11): 2187-2191
- Hande KR, Wedlund PJ, Noone RM, et al. Pharmacokinetics of high-dose etoposide (VP-16-213) administered to cancer patients. *Cancer Res* 1984; 44: 379-382
- Hansen HH, Muggia FM. Treatment of malignant brain tumors with nitrosoureas. *Cancer Chemother Rep* 1971; 55 (2): 99-100
- Harisiadis L, Chang CH. Medulloblastoma in children: a correlation between staging and results of treatment. *Int J Radiat Oncol Biol Phys* 1977; 2: 833-841
- Henwood JM, Brogden RN. Etoposide. A review of its pharmacodynamic and pharmacokinetic properties, and therapeutic potential in combination chemotherapy of cancer. *Drugs* 1990; 39: 438-490
- Hodgson DC, Goumnerova LC, Loeffler JS, et al. Radiosurgery in the management of pediatric brain tumors. *Int J Radiat Oncol Biol Phys* 2001; 50 (4): 929-935
- Hongeng S, Visudtibhan A, Dhanachai M, et al. Treatment of leptomeningeal relapse of medulloblastoma with temozolomide. *J Pediatr Hematol Oncol* 2002; 24 (7): 591-593
- Hughes EN, Shillito J, Sallan SE, et al. Medulloblastoma at the Joint Center for Radiation Therapy between 1968 and 1984. *Cancer* 1988; 61: 1992-1998
- Hughes WT; Armstrong D, Bodey GP, et al. 2002 Guidelines for the use of antimicrobial agents in neutropenic patients with cancer. *Clin Infect Dis* 2002; 34: 730-751
- Jaeckle KA, Hess KR, Yung WK, et al. North American Brain Tumor Consortium. Phase II evaluation of temozolomide and 13-cis-retinoic acid for the treatment of recurrent and progressive malignant glioma: a North American Brain Tumor Consortium study. *J Clin Oncol* 2003; 21 (12): 2305-2311
- Jennison, C, Turnbull B. Group sequential methods with applications to clinical trials. London, Chapman & Hall. 2000; chapter 5.2: 111-123

- Kaatsch P, Rickert CH, Kühl J, et al. Population-based epidemiologic data on brain tumors in German children. *Cancer* 2001; 92 (12): 3155-3164
- Kalifa C, Flamant F, Hartmann O, et al. Postsurgical chemotherapy without radiation therapy in patients less than three years with malignant brain tumors. *Pediatr Neurosci* 1988; 14: 163 (abstr)
- Kalifa C, Valteau D, Pizer B, et al. High-dose chemotherapy in childhood brain tumours. *Childs Nerv Syst* 1999; 15: 498-505
- Kellie SJ. Chemotherapy of central nervous system tumours in infants. *Childs Nerv Syst* 1999; 15 (10): 592-612
- Kennedy C, Calaminus G. A comparison between two quality of life measures in survivors of brain tumors or leukaemia in childhood and adolescents. *Med Pediatr Oncol* 2002; 39: 345
- Kim S, Chatelut E, Kim JC, et al. Extended CSF cytarabine exposure following intrathecal administration of DTC 101. *J Clin Oncol* 1993; 11 (11): 2186-2193
- Kiya K, Uozumi T, Ogasawara H, et al. Penetration of etoposide into human malignant brain tumors after intravenous and oral administration. *Cancer Chemother Pharmacol* 1992; 29: 339-342
- Kleihues P, Cavenee WK. World Health Organization Classification of Tumours. Pathology Genetics. Tumours of the Nervous System. Lyon: IARC Press; 2000
- Kleiman G, Hochberg F, Richardson E. Systemic metastases from medulloblastoma: Report of two cases and review of the literature. *Cancer* 1984; 48: 2296-2309
- Kochi M, Kuratsu JI, Mihara Y, et al. Ventriculolumbar perfusion of 3-[(4-amino-2-methyl-5-pyrimidinyl)methyl]-1-(2-chlorethyl)-1-nitrosurea hydrochloride. *Neurosurgery* 1993; 33 (5): 817-823
- Kortmann RD, Timmermann B, Becker G, et al. Advances in treatment techniques and time/dose schedules in external radiation therapy of brain tumours in childhood. *Klin Pädiatr* 1998; 210: 220-226
- Kortmann RD, Becker G, Perelmouter J, et al. Geometric accuracy of field alignment in fractionated stereotactic conformal radiotherapy of brain tumors. *Int J Radiat Oncol Biol Phys* 1999; 43 (4): 921-926
- Kortmann RD, Timmermann B, Kühl J, et al. HIT'91 (prospective co-operative study for the treatment of malignant brain tumours in childhood): accuracy and acute toxicity of the irradiation of the craniospinal axis. *Strahlenther Onkol* 1999; 175 (4): 162-169
- Kortmann, RD, Kühl J, Timmermann B, et al. Postoperative neoadjuvant chemotherapy before radiotherapy as compared to immediate radiotherapy followed by maintenance chemotherapy in the treatment of medulloblastoma in childhood: results of the German prospective randomized trial HIT'91. *Int J Radiat Oncol Biol Phys* 2000; 46 (2): 269-279
- Kosch A, Mölenkamp G, Däumling E, et al. Assessment of independence in daily life in pediatric oncology by FMH-questionnaire. *Klin Pädiatr* 1998; 210(6): 390-394
- Kovalic JJ, Flaris N, Grigsby PW, et al. Intracranial ependymoma long term outcome, patterns of failure. *J Neurooncol* 1993; 15: 125-131
- Kovnar E, Kun L, Krischer J. Patterns of dissemination and recurrence in childhood ependymoma: preliminary results of Pediatric Oncology Group protocol #8532. *Ann Neurol* 1991; 30: 457 (abstr)
- Kühl J, Müller HL, Berthold F, et al. Preradiation chemotherapy of children and young adults with malignant brain tumors: results of the German pilot trial HIT'88/89. *Klin Pädiatr* 1998; 210: 227-233
- Kühl J. Protokoll HIT 2000, Version vom 01.06.2000
- Kun LE, Kovnar EH, Sanford RA. Ependymomas in children. *Pediatr Neurosci* 1988; 14: 57-63
- Lefkowitz IB, Packer RJ, Ryan SG, et al. Late recurrence of primitive neuroectodermal tumor/medulloblastoma. *Cancer* 1988; 62: 826-830
- Lefkowitz JB, Packer RJ, Siegel KR, et al. Results of treatment of children with recurrent primitive neuroectodermal tumors-medulloblastoma with lomustine, cisplatin and vincristine. *Cancer* 1990; 65: 412-417
- Mahoney DHJ, Strother D, Camitta B, et al. High-dose melphalan and cyclophosphamide with autologous bone marrow rescue for recurrent/progressive malignant brain tumors in children: a pilot pediatric oncology group study. *J Clin Oncol* 1996; 14: 382-388
- Mason WP, Goldman S, Yates AJ, et al. Survival following intensive chemotherapy with bone marrow reconstitution for children with recurrent intracranial ependymoma--a report of the Children's Cancer Group. *J Neurooncol* 1998; 37 (2): 135-143

- McIntosh N. Medulloblastoma: a changing prognosis. *Arch Dis Child* 1979; 54: 200-203
- Michon J, Schleiermacher G. Autologous haematopoietic stem cell transplantation for pediatric solid tumors. *Baillieres Best Pract Res Clin Haematol* 1999; 12: 247-259
- Milker-Zabel S, Zabel A, Thilmann C, et al. Results of three-dimensional stereotactically-guided radiotherapy in recurrent medulloblastoma. *J Neurooncol* 2002; 60 (3): 227-233
- Müller HL, Fröhwald MC, Scheubeck M, et al. A possible role for somatostatin receptor scintigraphy in the diagnosis and follow-up of children with medulloblastoma. *J Neurooncol*. 1998; 38 (1): 27-40
- Müller HL, Bueb K, Bartels U, et al. Obesity after childhood craniopharyngioma - German multicenter study on preoperative risk factors and quality of life. *Klin Pädiatr* 2001; 213(4): 244-249
- Needle MN, Goldwein JW, Grass J, et al. Adjuvant chemotherapy for the treatment of intracranial ependymoma of childhood. *Cancer* 1997a; 80: 341-347
- Needle MN, Molloy PT, Geyer JR, et al. Phase II study of daily oral etoposide in children with recurrent brain tumors and other solid tumors. *Med Pediatr Oncol* 1997b; 29 (1): 28-32
- Newton HB, Henson J, Walker RW. Extraneural metastases in ependymoma. *J Neurooncol* 1992; 14: 135-142
- nQuery Advisor, Version 3.0. Statistical Solutions Ltd., Cork, Ireland.
- O'Brien P, Fleming T. A multiple testing procedure for clinical trials. *Biometrics* 1979; 35: 549-556
- Oken, M.M., Creech, R.H., Tormey, D.C., Horton, J., Davis, T.E., McFadden, E.T., Carbone, P.P.: Toxicity And Response Criteria Of The Eastern Cooperative Oncology Group. *Am J Clin Oncol* 5:649-655, 1982.
- O'Reilly SM, Newlands ES, Glaser MG, et al. Temozolomide: a new oral cytotoxic chemotherapeutic agent with promising activity against primary brain tumours. *Eur J Cancer* 1993; 29° (7): 940-942
- Packer RJ, Sutton LN, Eltermann R, et al. Outcome for children with medulloblastoma treated with radiation and cisplatin, CCNU, and vincristine chemotherapy. *J Neurosurg* 1994; 81: 690-698
- Patrice SJ, Tarbell NJ, Goumnerova LC, et al. Results of radiosurgery in the management of recurrent and residual medulloblastoma. *Pediatr Neurosurg* 1995; 22 (4): 197-203
- Pendergrass TW, Milstein JM, Geyer JR, et al. Eight drugs in one day chemotherapy for brain tumors: Experience in 107 children and rationale for preradiation chemotherapy. *J Clin Oncol* 1987; 5: 1221-1231
- Postmus PE, Holthuis JJ, Haaxma-Reiche H, et al. Penetration of VP 16-213 into cerebrospinal fluid after high-dose intravenous administration. *J Clin Oncol* 1984; 2: 215-220
- Riccardi R, Cefalo G, Ruggiero A, et al. High response rate to temozolomide in heavily pretreated children with medulloblastoma. *Proceedings of 93<sup>rd</sup> Annual Meeting AACR* 2002, 43: abstr 3711
- Rosen G, Ghavimi F, Nirenberg A, et al. High-dose methotrexate with citrovorum factor rescue for the treatment of central nervous system tumors in children. *Cancer Treat Rep* 1977; 61: 781-790
- Rousseau P, Habrand JL, Sarrazin D, et al. Treatment of intracranial ependymomas of children: review of 15-year experience. *J Radiat Oncol Biol Phys* 1994; 28: 381-386
- Rubinstein LJ. Justification for a cytogenetic scheme of embryonal central neuroepithelial tumors. In: Fields WS (ed). Primary brain tumors. A review of histologic classification. Springer-Verlag, New York, 1989: 16-27
- Rutkowski S, Bode U, Deinlein F, et al. Cure of children less than three years with medulloblastoma (M0/M1-stage) by postoperative chemotherapy only: final results of the HIT-SKK'92 Study. 11<sup>th</sup> International Symposium on Pediatric Neuro-Oncology, Boston, June 13<sup>th</sup>-16<sup>th</sup>, 2004 (abstr)
- Rutkowski S, Bode U, Deinlein F, et al. Treatment of early childhood medulloblastoma by postoperative chemotherapy alone. *N Engl J Med*. 2005; 352(10): 978-986
- Santana VM, Schell MJ, Williams R, et al. Escalating sequential high-dose carboplatin and etoposide with autologous marrow support in children with relapsed solid tumors. *Bone Marrow Transplant* 1992; 10: 457-462
- Savaraj N, Lu K, Feun LG, et al. Comparison of CNS penetration, tissue distribution, and pharmacology of VP 16-213 by intracarotid and intravenous administration in dogs. *Cancer Invest* 1987; 5: 11-16
- Schäfer H, Müller HH. Modification of the sample size and the schedule of interim analyses in survival trials based on data inspections. *Statistics in Medicine* 2001; 20: 3741-3751
- Scheurlen WG, Schwabe GC, Joos S, et al. Molecular analysis of childhood primitive neuroectodermal tumors defines markers associated with poor outcome. *J Clin Oncol* 1998; 16 (7): 2478-2485

- Segal RA, Goumnerova LC, Kwon YK, et al. Expression of the neurotrophin receptor TrkC is linked to a favorable outcome in medulloblastoma. *Proc Natl Acad Sci USA* 1994; 91 (26): 12867-12871
- Shapiro WR. Chemotherapy of primary malignant brain tumors in children. *Cancer* 1975; 35(3 suppl): 965-972
- Shepherd SF, Laing RW, Cosgrove VP, et al. Hypofractionated stereotactic radiotherapy in the management of recurrent glioma. *Int J Radiat Oncol Biol Phys* 1997; 37 (2): 393-398
- Simon A, Fleischhack G. Nicht medikamentöse Infektionsprophylaxe in der pädiatrischen Hämatologie-Onkologie. *Klin Pädiatr* 2001; 213 Sonderheft 1: A9-A21
- Simon A, Fleischhack G, Marklein G, et al. Chemoprophylaxe bakterieller Infektionskrankheiten in der pädiatrischen Onkologie. *Klin Pädiatr* 2001; 213 Sonderheft 1: A22-A37
- Simon R. Optimal two stage designs for phase II clinical trials. *Controlled Clinical Trials* 1989; 10: 1-10
- Stafford SL, Pollock BE, Foote RL, et al. Stereotactic radiosurgery for recurrent ependymoma. *Cancer* 2000; 88 (4): 870-875
- Sterba J, Pavelka Z, Slampa P. Concomitant radiotherapy and metronomic temozolomide in pediatric high-risk brain tumors. *Neoplasma* 2002; 49 (2): 117-120
- Stremetzne S, Jaehde U, Schunack W. Determination of the cytotoxic catechol metabolite of etoposide (3'-O-demethyletoposide) in human plasma by high-performance liquid chromatography. *J Chromatogr B Biomed Sci Appl* 1997; 703: 209-215
- Strother D, Ashley D, Kellie SJ, et al. Feasibility of four consecutive high-dose chemotherapy cycles with stem-cell rescue for patients with newly diagnosed medulloblastoma or supratentorial primitive neuroectodermal tumor after craniospinal radiotherapy: results of a collaborative study. *J Clin Oncol* 2001; 19 (10): 2696-2704
- Suh JH, Barnett GH. Stereotactic radiosurgery for brain tumors in pediatric patients. *Technol Cancer Res Treat* 2003; 2 (2): 141-146
- Sur P, Sribnick EA, Patel SJ, Ray SK, Banik NL. Dexamethasone decreases temozolomide-induced apoptosis in human glioblastoma T98G cells. *Glia* 2005; 50(2): 160-167
- Tait DM, Thorton-Jones H, Bloom HJG, et al. Adjuvant chemotherapy for medulloblastoma: The first multi-centre control trial of the International Society of Paediatric Oncology (SIOP I). *Eur J Cancer* 1990; 26: 464-447
- Taylor RE. Review of radiotherapy dose and volume for intracranial ependymoma. *Pediatr Blood Cancer* 2004; 42 (5): 457-460
- Timmermann B, Kortmann RD, Kühl J, et al. Combined postoperative irradiation and chemotherapy for anaplastic ependymomas in childhood: results of the German prospective trials HIT 88/89 and HIT 91. *Int J Radiat Oncol Biol Phys* 2000; 46: 287-295
- Timmermann B, Kortmann RD, Kühl J, et al. Postoperative therapy of ependymoma in children: results of the German HIT-SKK and HIT trials. *Med Pediatr Oncol* 2002; 39: 249
- Tomlinson FH, Lihou MG, Smith PJ. Comparison of in vitro activity of epipodophyllotoxins with other chemotherapeutic agents in human medulloblastomas. *Br J Cancer* 1991; 64: 1051-1059
- Torres CF, Rebsamen S, Silber JH, et al. Surveillance scanning of children with medulloblastoma. *N Engl J Med* 1994; 330: 892-895
- Trimis G, Prodromou F, Sfakianos G, et al. Temozolomide in malignant CNS Neoplasms of Childhood: a single institution experience. *Pediatr Blood Cancer* 2004, 43 (4): 401 (abstr)
- van der Gaast A, Sonneveld P, Mans DR, et al. Intrathecal administration of etoposide in the treatment of malignant meningitis: feasibility and pharmacokinetic data. *Cancer Chemother Pharmacol* 1992; 29: 335-337
- van Veelen-Vincent ML, Pierre-Kahn A, Kalifa C, et al. Ependymoma in childhood: prognostic factors, extent of surgery, and adjuvant therapy. *J Neurosurg* 2002; 97: 827-835
- Vera K, Djafari L, Faivre S, et al. Dose-dense regimen of temozolomide given every other week in patients with primary central nervous system tumors. *Ann Oncol* 2004; 15 (1): 161-171
- Verschraegen CF, Jaeckle K, Giovanella B, et al. Alternative administration of camptothecin analogues. *Ann N Y Acad Sci* 2000; 922: 237-246
- Wagner S, Erdlenbruch B, Längler A, et al. Oral topotecan in children with recurrent or progressive high-grade glioma: a Phase I/II study by the German Society for Pediatric Oncology and Hematology. *Cancer* 2004; 100 (8): 1750-1757
- Wald, A. Sequential tests of statistical hypotheses. *Annals of Mathematical Statistics* 1945; 16: 117-186.

Walker RW, Allen JC. Cisplatin in the treatment of recurrent childhood primary brain tumors. *J Clin Oncol* 1988; 6: 62-66

Warmuth-Metz M, Kühl J, Bendzus M., et al. Die Inzidenz von Marklagerveränderungen bei hochdosierter Methotrexat-Gabe bei Kindern mit Medulloblastomen unter 3 Jahren: Vorläufige Ergebnisse der HIT-SKK-Studien. *Aktuelle Neuropädiatrie* 1998, Herausgeber: Bentele und Kohlschütter, 1998: 77-82

Wassmer G, Eisebitt R. Institut for Medical Statistics, University of Cologne, Clin Research GmbH, Cologne, Version 2.1.4, January 2004.

Wolff JE, Molenkamp G, Westphal S, et al. Oral trofosfamide and etoposide in pediatric patients with glioblastoma multiforme. *Cancer* 2000; 89 (10): 2131-2137

Wolff JE, Westphal S, Molenkamp G, et al. Treatment of paediatric pontine glioma with oral trophosphamide and etoposide. *Br J Cancer* 2002; 87 (9): 945-949

Woo C, Stea B, Lulu B, Hamilton A, Cassady JR. The use of stereotactic radiosurgical boost in the treatment of medulloblastomas. *Int J Radiat Oncol Biol Phys* 1997; 37 (4): 761-764

Young JL, Miller W. Incidence of malignant tumors in U.S. children. *J Pediatr* 1975; 86: 254-258

Zernikow B, Griessinger N, Fengler R. [Practice of pain control in paediatric oncology--recommendations of the quality-monitoring group of the German Society for Paediatric Oncology and Haematology (GPOH)] *Schmerz* 1999; 13(3): 213-235

Zia MI, Forsyth P, Chaudhry A, et al. Possible benefits of high-dose chemotherapy and autologous stem cell transplantation for adults with recurrent medulloblastoma. *Bone Marrow Transplant* 2002; 30 (9): 565-569

Ziegler A, Lange S, Bender R. Überlebenszeitanalyse: Der Log-Rang-Test. *Dtsch Med Wochenschr* 2004; 129: T4-T6